

COMPARISON OF MACNEILLE, CANONICAL, AND PROFINITE COMPLETIONS

GURAM BEZHANISHVILI AND JACOB VOSMAER

ABSTRACT. Using duality theory, we give necessary and sufficient conditions for the MacNeille, canonical, and profinite completions of distributive lattices, Heyting algebras, and Boolean algebras to be isomorphic.

1. INTRODUCTION

In the theory of lattice completions and in applications of lattice theory to logic, the MacNeille and canonical completions play a fundamental role. The MacNeille completions provide completeness of various predicate logics with respect to their algebraic semantics, as was shown in [19, 16, 17]. On the other hand, the Jónsson–Tarski representation theorem for Boolean algebras with operators [14] is the key for many completeness results in non-classical propositional logics (such as modal and superintuitionistic logics). A connection between the MacNeille and canonical completions was discussed in [9]. Yet another completion—the profinite completion—has its origins in Galois theory. It has recently been shown that there is a close connection between the canonical and profinite completions [3, 12]. The aim of this paper is to compare these three completions for distributive lattices, Heyting algebras, and Boolean algebras. Our main tool is duality theory.

We give a necessary and sufficient condition for the MacNeille completion \bar{L} of a bounded distributive lattice L to be isomorphic to its canonical completion L^σ . Since L^σ is isomorphic to the profinite completion \hat{L} of L [3, Thm. 2.11], we obtain a criterion for the three completions to be isomorphic. The isomorphisms, however, may not commute with the embeddings $\eta : L \hookrightarrow \bar{L}$, $\zeta : L \hookrightarrow L^\sigma$, and $\iota : L \hookrightarrow \hat{L}$. We give necessary and sufficient conditions for the isomorphisms to commute with η , ζ , and ι . When L happens to be a Heyting algebra, our results about the isomorphism between \bar{L} and L^σ apply unchanged. But now, unlike the case of bounded distributive lattices, L^σ is not necessarily isomorphic to \hat{L} . Therefore, we add necessary and sufficient conditions for the existence of an isomorphism between L^σ and \hat{L} , and between \bar{L} and \hat{L} . Consequently, we obtain necessary and sufficient conditions for the three completions of a Heyting algebra to be isomorphic. We also give necessary and sufficient conditions for these isomorphisms to commute with η , ζ , and ι .

The paper is organized as follows. In Section 2 we recall the definitions and basic facts about the MacNeille, canonical, and profinite completions. In Section 3 we recall the Priestley duality for bounded distributive lattices, and then establish our first main results about the isomorphism of the MacNeille, canonical, and profinite completions of bounded distributive lattices. In Section 4 we briefly recall the Esakia duality for Heyting algebras, and then prove

2000 *Mathematics Subject Classification.* 06B23; 06D20.

Key words and phrases. MacNeille completion, canonical completion, profinite completion.

The second author was supported by VICI grant 639.073.501 of the Netherlands Organization for Scientific Research (NWO).

our main results about the isomorphism of the three completions for Heyting algebras. We also discuss consequences of our results for the case of Boolean algebras. Finally, in Section 5 we briefly discuss extensions of our results to more general settings.

2. PRELIMINARIES

In this paper we are interested in the conditions under which different completions of distributive lattices and related algebras are isomorphic. We are mostly concerned with the following three completions: the MacNeille completion, the canonical completion, and the profinite completion. In this preliminary section we introduce the main definitions from the theory of completions of lattices to make the paper self-contained.

Let P be a poset. For $a \in P$ let $\uparrow a = \{x \in P \mid a \leq x\}$ and $\downarrow a = \{x \in P \mid x \leq a\}$. We recall that $S \subseteq P$ is *join-dense* in P if for all $a \in P$ we have $\bigvee(\downarrow a \cap S)$ exists and $a = \bigvee(\downarrow a \cap S)$. The notion of a *meet-dense* subset of P is dual. The next lemma is well-known (see, e.g., [6, p. 160]).

Lemma 2.1. *Let L be a lattice and let $S \subseteq L$.*

- (1) *S is join-dense in L iff for each $a, b \in L$ with $a \not\leq b$, there exists $s \in S$ such that $s \leq a$ and $s \not\leq b$.*
- (2) *S is meet-dense in L iff for each $a, b \in L$ with $a \not\leq b$, there exists $s \in S$ such that $a \not\leq s$ and $b \leq s$.*

Throughout we assume that **all lattices are bounded**. If not, then we can always add a new top and bottom to them. The next definition is well-known; see, e.g., [21, 15]. Note that we do *not* assume that L is a complete lattice.

Definition 2.2. *Let L be a lattice.*

- (1) *For $p, q \in L$ we say that a pair (p, q) splits L if $\uparrow p \cap \downarrow q = \emptyset$ and $\uparrow p \cup \downarrow q = L$. When (p, q) splits L , then we call (p, q) a splitting pair, p a splitting element, and q a co-splitting element. Let $S(L)$ denote the set of splitting elements of L and $CS(L)$ denote the set of co-splitting elements of L .*
- (2) *We say that $0 \neq p \in L$ is join-prime if from $p \leq a \vee b$ it follows that $p \leq a$ or $p \leq b$. The notion of a meet-prime element is dual. Let $J(L)$ and $M(L)$ denote the sets of join-prime and meet-prime elements of L , respectively.*
- (3) *We say that $0 \neq p \in L$ is completely join-prime if for each $S \subseteq L$ such that $\bigvee S$ exists and $p \leq \bigvee S$, there exists $s \in S$ with $p \leq s$. The notion of a completely meet-prime element is dual. Let $J^\infty(L)$ and $M^\infty(L)$ denote the sets of completely join-prime and completely meet-prime elements of L , respectively.*

It is easy to see that $S(L) \subseteq J^\infty(L) \subseteq J(L)$ and that $CS(L) \subseteq M^\infty(L) \subseteq M(L)$. Moreover, if L is complete, then $S(L) = J^\infty(L)$ and $CS(L) = M^\infty(L)$. To see this, observe that from $p \in J^\infty(L)$ (resp. $q \in M^\infty(L)$) it follows that $(p, \bigvee\{x \in L \mid p \not\leq x\})$ (resp. $(\bigwedge\{x \in L \mid x \not\leq q\}, q)$) is a splitting pair, and so $p \in S(L)$ (resp. $q \in CS(L)$). Furthermore, if L is finite, then it is obvious that in addition we have $J^\infty(L) = J(L)$ and $M^\infty(L) = M(L)$. An example of a complete lattice in which $J^\infty(L) \subset J(L)$ (resp. $M^\infty(L) \subset M(L)$) is the lattice of closed (resp. open) subsets of the real line. Indeed, singletons (resp. complements of singletons) are join-prime (resp. meet-prime), but are not completely join-prime (resp. completely meet-prime). An example of an incomplete lattice in which $S(L) \subset J^\infty(L)$ (resp. $CS(L) \subset M^\infty(L)$) is given below.




FIGURE 1. $(\omega \times \omega) \oplus \omega^{\text{op}}$

Example 2.3. Let ω be the first infinite ordinal with its usual order, and consider the (distributive) lattice $L = (\omega \times \omega) \oplus \omega^{\text{op}}$, where \oplus is the operation of sum and ω^{op} is the opposite of ω (see Fig. 1). Clearly L is not complete. Moreover, each $p = (0, n)$ or $(n, 0)$ is in $J^\infty(L)$, but does not belong to $S(L)$. By taking the opposite of L we obtain a lattice in which not every completely meet-prime element is co-splitting.

Definition 2.4. Let L be a lattice.

- (1) [5, p. 211] We call L *principally separated* if for each $a, b \in L$ with $a \not\leq b$, there is a *splitting pair* (p, q) such that $p \leq a$ and $b \leq q$.
- (2) We call L *completely join-prime generated* if $J^\infty(L)$ is *join-dense* in L .

Remark 2.5. Using Lemma 2.1, it is easy to see that if L is principally separated, then $S(L)$ is join-dense in L . Since $S(L) \subseteq J^\infty(L)$, it follows that each principally separated lattice is completely join-prime generated. On the other hand, the lattice given in Example 2.3 is an example of a completely join-prime generated lattice, which is not principally separated. Nevertheless, the observations in the paragraph following Definition 2.2 imply that the two notions are equivalent in the setting of complete lattices.

Definition 2.6. Let L be a lattice. A *completion* of L is a pair (C, ξ) where C is a complete lattice and $\xi: L \rightarrow C$ is a lattice embedding. Given two completions of L , (C, ξ) and (D, μ) , and a lattice isomorphism $f: C \rightarrow D$, we call $f: (C, \xi) \rightarrow (D, \mu)$ an *isomorphism of completions* if $f \circ \xi = \mu$. We say (C, ξ) and (D, μ) are *isomorphic as completions of L* if there exists an isomorphism of completions $f: (C, \xi) \rightarrow (D, \mu)$.

Now we recall the definitions of the MacNeille and canonical completions of lattices.

Definition 2.7. Let L be a lattice.

- (1) [1] *The MacNeille completion of L is a pair (\bar{L}, η) , where \bar{L} is a complete lattice and $\eta: L \rightarrow \bar{L}$ is a lattice embedding such that $\eta[L]$ is both join-dense and meet-dense in \bar{L} .*
- (2) [8] *The canonical completion of L is a pair (L^σ, ζ) , where L^σ is a complete lattice and $\zeta: L \rightarrow L^\sigma$ is a lattice embedding such that:

 - (i) *For each filter F and ideal I of L , from $F \cap I = \emptyset$ it follows that $\bigwedge \zeta[F] \not\leq \bigvee \zeta[I]$.*
 - (ii) *The set $K_L = \{\bigwedge \zeta[S] \mid S \subseteq L\}$ of closed elements is join-dense in L^σ .*
 - (iii) *The set $O_L = \{\bigvee \zeta[S] \mid S \subseteq L\}$ of open elements is meet-dense in L^σ .**

The third completion we will be dealing with in this paper is the profinite completion. Let L be a lattice and let $\Phi_L = \{\theta \in \text{Con } L \mid L/\theta \text{ is finite}\}$ be the congruences of L of finite index. For $\theta, \psi \in \Phi_L$ with $\theta \subseteq \psi$, let $h_{\theta\psi}: L/\theta \rightarrow L/\psi$ be given by $h_{\theta\psi}(a/\theta) = a/\psi$. Then (Φ_L, \supseteq) is a directed poset, and so $\langle \{L/\theta\}, h_{\theta\psi} \rangle$ is an inverse system of finite homomorphic images of L . We denote the inverse limit of $\langle \{L/\theta\}, h_{\theta\psi} \rangle$ by \widehat{L} . It is well-known that

$$\widehat{L} = \{\alpha \in \prod_{\Phi_L} L/\theta \mid \forall \theta, \psi \in \Phi_L \text{ with } \theta \subseteq \psi, \text{ if } \alpha(\theta) = a/\theta \text{ then } \alpha(\psi) = a/\psi\}.$$

We define $\iota: L \rightarrow \widehat{L}$ by $\iota(a) = \alpha_a$, where $\alpha_a(\theta) = a/\theta$ for $\theta \in \Phi_L$. We note that $\iota: L \rightarrow \widehat{L}$ may not be 1-1 in general, but ι is 1-1 if L is distributive (see, e.g., [3, pp. 145–146]).

Definition 2.8. [3] *Let L be a distributive lattice. The profinite completion of L is the pair (\widehat{L}, ι) .*

In this paper we are primarily concerned with two questions. Firstly, we want to find necessary and sufficient conditions for the lattices \bar{L} , L^σ , and \widehat{L} to be isomorphic. Secondly, we seek necessary and sufficient conditions for the completions (\bar{L}, η) , (L^σ, ζ) , and (\widehat{L}, ι) to be isomorphic as completions of L .

3. DISTRIBUTIVE LATTICES

We briefly recall the basics of Priestley duality [18] for distributive lattices. For a topological space X , let \mathbf{I} and \mathbf{C} denote the interior and closure operators of X , respectively. We recall that a subset U of X is *clopen* if U is both closed and open, and that X is *zero-dimensional* if clopen subsets of X form a basis for the topology. We also recall that X is a *Stone space* if X is compact, Hausdorff, and zero-dimensional.

Let $\langle X, \leq \rangle$ be a partially ordered set. We call $A \subseteq X$ an *upset* of X if $x \in A$ and $x \leq y$ imply $y \in A$. Let $\text{Up}(X)$ denote the set of upsets of X .

Definition 3.1. *A Priestley space is a triple $\langle X, \tau, \leq \rangle$, where $\langle X, \tau \rangle$ is a compact space, $\langle X, \leq \rangle$ is a partially ordered set, and $\langle X, \tau, \leq \rangle$ satisfies the Priestley separation axiom: For $x, y \in X$ with $x \not\leq y$, there exists a clopen upset U of X such that $x \in U$ and $y \notin U$. (Note that the above two conditions imply that $\langle X, \tau \rangle$ is a Stone space.)*

By Priestley duality, the dual of a distributive lattice L is the Priestley space $\langle \text{Pr}(L), \tau_L, \subseteq \rangle$, where $\text{Pr}(L)$ is the set of prime filters of L and τ_L is the topology generated by $\{\varphi(a) - \varphi(b) \mid a, b \in L\}$; here $\varphi(a) = \{\nabla \in \text{Pr}(L) \mid a \in \nabla\}$. Conversely, the dual of a Priestley space $\langle X, \tau, \leq \rangle$ is the distributive lattice $\langle \text{Up}_\tau(X), \cap, \cup, \emptyset, X \rangle$, where $\text{Up}_\tau(X)$ is the set of clopen upsets of X . Moreover, the function $\varphi: L \rightarrow \text{Up}_{\tau_L}(\text{Pr}(L))$ is a lattice isomorphism. Thus, each distributive lattice L is represented as the lattice $\text{Up}_\tau(X)$ of clopen upsets of some Priestley space X .

Let $\langle X, \tau, \leq \rangle$ be a Priestley space. If $S \subseteq X$, we write $\downarrow S = \{x \in X \mid \exists s \in S : x \leq s\}$ ($\uparrow S$ is defined dually). Following [13, Section 3], we define the operators \mathbf{J} and \mathbf{D} on the powerset of X as follows:

$$(1) \quad \mathbf{J}S = -\downarrow - \mathbf{I}S,$$

$$(2) \quad \mathbf{D}S = \uparrow \mathbf{C}S.$$

Then for each $S \subseteq X$ we have $\mathbf{J}S$ is the largest open upset contained in S and $\mathbf{D}S$ is the smallest closed upset containing S . Now define

$$\mathcal{RO}(X) = \{S \subseteq X \mid \mathbf{JDS} = S\}.$$

Because \mathbf{J} and \mathbf{D} form a Galois connection between the complete lattices of open upsets and closed upsets of X , we have that $\mathcal{RO}(X)$ is a complete lattice under set-theoretic inclusion [13, Lemma 3.4].

Lemma 3.2. *Let L be a distributive lattice and let $\langle X, \tau, \leq \rangle$ be its Priestley space. Then:*

- (1) *There is an isomorphism $f : \bar{L} \rightarrow \mathcal{RO}(X)$ such that $f \circ \eta = \varphi$. Therefore, $(\bar{L}, \eta) \simeq (\mathcal{RO}(X), \varphi)$.*
- (2) *There is an isomorphism $g : L^\sigma \rightarrow \text{Up}(X)$ such that $g \circ \zeta = \varphi$. Therefore, $(L^\sigma, \zeta) \simeq (\text{Up}(X), \varphi)$.*
- (3) *There is an isomorphism $h : \hat{L} \rightarrow \text{Up}(X)$ such that $h \circ \iota = \varphi$. Therefore, $(\hat{L}, \iota) \simeq (\text{Up}(X), \varphi)$.*

Proof. (1) As in [13], we view \bar{L} , up to isomorphism, as the lattice of normal ideals of L . Then $\eta : L \rightarrow \bar{L}$ is given by $\eta(a) = \downarrow a$. It was shown in [13, Theorem 3.5] that $f : \bar{L} \rightarrow \mathcal{RO}(X)$, given by $f(I) = \bigcup \{\varphi(a) \mid a \in I\}$, is a lattice isomorphism. We show that $f \circ \eta = \varphi$. For $a \in L$ we have $f(\eta(a)) = f(\downarrow a) = \bigcup \{\varphi(b) \mid b \leq a\} = \varphi(a)$.

(2) is well-known; see, e.g., [10, Section 2].

(3) It was shown in [3, Theorem 2.11] that L^σ is isomorphic to \hat{L} . Consequently, \hat{L} is isomorphic to $\text{Up}(X)$. The isomorphism $h : \hat{L} \rightarrow \text{Up}(X)$ can be defined explicitly as follows. Let $\theta \in \Phi_L$. It is well-known that the 1-1 correspondence between congruences of L and closed subsets of X restricts to the 1-1 correspondence between elements of Φ_L and finite subsets of X . Let X_θ be the finite subset of X corresponding to θ . For $\alpha \in \hat{L}$ we have $\alpha(\theta) = a/\theta$ for some $a \in L$. Then $\varphi(a) \cap X_\theta$ is an upset of X_θ . If there is another $b \in L$ such that $\alpha(\theta) = b/\theta$, then $\varphi(a) \cap X_\theta = \varphi(b) \cap X_\theta$. Thus, $\varphi(a) \cap X_\theta$ is independent of $a \in L$, and we denote it by $\varphi(\alpha(\theta))$. Let $U_\alpha = \bigcup \{\varphi(\alpha(\theta)) \mid \theta \in \Phi_L\}$. We show that U_α is an upset of X . If $x \in U_\alpha$ and $x \leq y$, then there is $\theta \in \Phi_L$ such that $x \in \varphi(\alpha(\theta))$. Since $\{x, y\}$ is a finite subset of X , there is $\kappa \in \Phi_L$ such that $X_\kappa = \{x, y\}$. Let $\rho = \theta \cap \kappa$. Then $X_\rho = X_\theta \cup X_\kappa$. Therefore, $y \in X_\rho \subseteq U_\alpha$, and so $U_\alpha \in \text{Up}(X)$. Define $h : \hat{L} \rightarrow \text{Up}(X)$ by $h(\alpha) = U_\alpha$. Then h is the desired isomorphism. We show that $h \circ \iota = \varphi$. For $a \in L$ we have $h(\iota(a)) = h(\alpha_a) = U_{\alpha_a}$. Since $U_{\alpha_a} = \bigcup \{\varphi(a) \cap X_\theta \mid \theta \in \Phi_L\}$, then U_{α_a} is the union of finite subsets of $\varphi(a)$. But so is $\varphi(a)$. Thus, $U_{\alpha_a} = \varphi(a)$, and so $h(\iota(a)) = \varphi(a)$. \square

We recall that a point x of a topological space X is *isolated* if $\{x\}$ is an open subset of X .

Definition 3.3. *Let $\langle X, \tau, \leq \rangle$ be a Priestley space. We call $x \in X$ an order-isolated point if both $\uparrow x$ and $\downarrow x$ are clopen. Let X_0 denote the set of order-isolated points of X .*

Since in a Priestley space $\langle X, \tau, \leq \rangle$ both $\uparrow x$ and $\downarrow x$ are always closed for each $x \in X$, we can equivalently define x to be order-isolated if both $\uparrow x$ and $\downarrow x$ are open. Clearly every order-isolated point is isolated, but the converse is not true as follows from [2, Example 2.8].

Theorem 3.4. *Let L be a distributive lattice and let X be its Priestley space.*

- (1) (p, q) splits L iff there is $x \in X_0$ such that $\varphi(p) = \uparrow x$ and $\varphi(q) = -\downarrow x$.
- (2) $p \in S(L)$ iff there is $x \in X_0$ such that $\varphi(p) = \uparrow x$.
- (3) $q \in CS(L)$ iff there is $x \in X_0$ such that $\varphi(q) = -\downarrow x$.
- (4) The posets $\langle S(L), \geq \rangle$, $\langle CS(L), \geq \rangle$, and $\langle X_0, \leq \rangle$ are order-isomorphic.

Proof. (1) First suppose that there is $x \in X_0$ such that $\varphi(p) = \uparrow x$ and $\varphi(q) = -\downarrow x$. Then $\varphi(p) \not\subseteq \varphi(q)$, and so $p \not\leq q$. Therefore, $\uparrow p \cap \downarrow q = \emptyset$. Moreover, for $a \in L$ we have that either $x \in \varphi(a)$ or $x \notin \varphi(a)$. If $x \in \varphi(a)$, then $\uparrow x \subseteq \varphi(a)$, as $\varphi(a)$ is an upset of X . Therefore, $\varphi(p) \subseteq \varphi(a)$, and so $a \in \uparrow p$. And if $x \notin \varphi(a)$, then $\downarrow x \cap \varphi(a) = \emptyset$. Thus, $\varphi(a) \subseteq -\downarrow x$, so $\varphi(a) \subseteq \varphi(q)$, and so $a \in \downarrow q$. Consequently, $\uparrow p \cup \downarrow q = L$, and so (p, q) splits L .

Conversely, suppose that (p, q) splits L . Then $p \in S(L)$ and $q \in CS(L)$. Therefore, p is join-prime and q is meet-prime. By [2, Thm. 2.7.1], there is $x \in X$ such that $\varphi(p) = \uparrow x$. A similar argument also gives us that there is $y \in X$ such that $\varphi(q) = -\downarrow y$. We show that $x = y$. Since $p \not\leq q$, we have $\varphi(p) \not\subseteq \varphi(q)$. Therefore, $\uparrow x \not\subseteq -\downarrow y$, so $\uparrow x \cap \downarrow y \neq \emptyset$, and so $x \leq y$. If $y \not\leq x$, then there is $a \in L$ such that $y \in \varphi(a)$ and $x \notin \varphi(a)$. It follows that $\uparrow x \not\subseteq \varphi(a)$, so $\varphi(p) \not\subseteq \varphi(a)$, and so $a \notin \uparrow p$. Since (p, q) splits L , we have $a \in \downarrow q$. Therefore, $\varphi(a) \subseteq \varphi(q) = -\downarrow y$. Thus, $\varphi(a) \cap \downarrow y = \emptyset$, and so $y \notin \varphi(a)$. The obtained contradiction proves that $y \leq x$, and so $x = y$. This implies that both $\uparrow x = \varphi(p)$ and $\downarrow x = -\varphi(q)$ are clopen, which means that $x \in X_0$. Consequently, there is $x \in X_0$ such that $\varphi(p) = \uparrow x$ and $\varphi(q) = -\downarrow x$.

(2) and (3) are immediate consequences of (1).

(4) It is obvious that $\langle S(L), \geq \rangle$ is isomorphic to $\langle CS(L), \geq \rangle$ (for $p \in S(L)$, there is a unique $q \in CS(L)$ such that (p, q) is a splitting pair, and sending p to q is the desired isomorphism). We show that $\langle CS(L), \geq \rangle$ is isomorphic to $\langle X_0, \leq \rangle$. Let $q \in CS(L)$. By (3), there is $x_q \in X_0$ such that $\varphi(q) = -\downarrow x_q$. We define $f: \langle CS(L), \geq \rangle \rightarrow \langle X_0, \leq \rangle$ by $f(q) = x_q$, and show that f is an order-isomorphism. That f is onto follows from (3). For $q, r \in CS(L)$, we have $q \geq r$ iff $\varphi(q) \supseteq \varphi(r)$ iff $-\downarrow x_q \supseteq -\downarrow x_r$ iff $\downarrow x_q \subseteq \downarrow x_r$ iff $x_q \leq x_r$ iff $f(q) \leq f(r)$, whence f is an order-isomorphism. \square

Theorem 3.5. *Let L be a distributive lattice and let X be its Priestley space. Then L is principally separated iff X_0 is dense in X .*

Proof. Suppose that L is principally separated and let $\varphi(a) - \varphi(b) \neq \emptyset$ be a basic open subset of X . From $\varphi(a) - \varphi(b) \neq \emptyset$ it follows that $a \not\leq b$. Therefore, there is a splitting pair (p, q) such that $p \leq a$ and $b \leq q$. By Theorem 3.4.1, there is $x \in X_0$ such that $\varphi(p) = \uparrow x$ and $\varphi(q) = -\downarrow x$. Thus, $x \in \varphi(a) - \varphi(b)$, whence X_0 is dense in X . Conversely, suppose that X_0 is dense in X . Let $a, b \in L$ with $a \not\leq b$. Then $\varphi(a) \not\subseteq \varphi(b)$, and so $\varphi(a) - \varphi(b)$ is a nonempty open subset of X . By density of X_0 , there is $x \in X_0 \cap (\varphi(a) - \varphi(b))$. Therefore, there are $p, q \in L$ such that $\varphi(p) = \uparrow x$ and $\varphi(q) = -\downarrow x$. By Theorem 3.4.1, (p, q) is a splitting pair. Moreover, $\varphi(p) \subseteq \varphi(a)$ and $\varphi(b) \subseteq \varphi(q)$. Therefore, $p \leq a$ and $b \leq q$, and so L is principally separated. \square

For a Priestley space $\langle X, \tau, \leq \rangle$, let X_{iso} denote the set of isolated points of X .

Lemma 3.6. *If X_0 is dense in X , then $X_0 = X_{\text{iso}}$.*

Proof. We already pointed out that X_0 is always a subset of X_{iso} . Suppose X_0 is dense in X and $x \in X_{\text{iso}}$. Then $\{x\} \cap X_0 \neq \emptyset$, implying that $x \in X_0$. Thus, $X_{\text{iso}} \subseteq X_0$. \square

Now we give necessary and sufficient conditions for the MacNeille completion of a distributive lattice to be isomorphic to the lattice of upsets of a poset. The equivalences (2) \Leftrightarrow (3) \Leftrightarrow (7) can be found in [5, Sec. 4] (see also [6, Sec. 4]). Our contribution is the dual characterization of such distributive lattices. Let L be a distributive lattice and let X be its Priestley space. For $a \in L$ we let $\varphi_0(a) = \varphi(a) \cap X_0$.

Theorem 3.7. *For a distributive lattice L and its Priestley space X , the following conditions are equivalent:*

- (1) \bar{L} is completely join-prime generated.
- (2) \bar{L} is principally separated.
- (3) L is principally separated.
- (4) X_0 is dense in X .
- (5) There exists an isomorphism of completions $\xi : (\bar{L}, \eta) \rightarrow (\text{Up}(X_0), \varphi_0)$.
- (6) $\bar{L} \simeq \text{Up}(X_0)$.
- (7) $\bar{L} \simeq \text{Up}(Y)$ for some poset Y .

Proof. The equivalence (1) \Leftrightarrow (2) is obvious since \bar{L} is complete; for (2) \Leftrightarrow (3) \Leftrightarrow (7) see [5, Sec. 4]; for (3) \Leftrightarrow (4) see Theorem 3.5; and (5) \Rightarrow (6) \Rightarrow (7) are immediate. Thus, to complete the proof, we need to show that (4) \Rightarrow (5).

(4) \Rightarrow (5): We define $\varepsilon : \mathcal{RO}(X) \rightarrow \text{Up}(X_0)$ by $\varepsilon(U) = U \cap X_0$. This function is well-defined because every $U \in \mathcal{RO}(X)$ is an upset. We show that ε is an order-isomorphism. For $U, V \in \mathcal{RO}(X)$ with $U \subseteq V$ we have $\varepsilon(U) = U \cap X_0 \subseteq V \cap X_0 = \varepsilon(V)$. Suppose that $U \not\subseteq V$. We show that $U \not\subseteq \mathbf{D}V$. If $U \subseteq \mathbf{D}V$, then $U \cap -\mathbf{D}V = \emptyset$. Since U is an upset, this implies $U \cap \downarrow -\mathbf{D}V = \emptyset$. Therefore, as $\mathbf{J}D\mathbf{V} = V$, we obtain $U \subseteq -\downarrow -\mathbf{D}V = \mathbf{J}D\mathbf{V} = V$, a contradiction. Thus, $U \not\subseteq \mathbf{D}V$, and so $U - \mathbf{D}V$ is a nonempty open subset of X . Because X_0 is dense in X , there exists $x \in X_0 \cap (U - \mathbf{D}V)$. This implies that $x \in U \cap X_0$ and $x \notin V \cap X_0$, and so $\varepsilon(U) \not\subseteq \varepsilon(V)$. Consequently, $U \subseteq V$ iff $\varepsilon(U) \subseteq \varepsilon(V)$. To see that ε is onto, let $U \in \text{Up}(X_0)$. We show that $\varepsilon(\mathbf{J}D\mathbf{U}) = U$. Clearly $U \subseteq \mathbf{J}D\mathbf{U} \cap X_0 = \varepsilon(\mathbf{J}D\mathbf{U})$. For the converse inclusion, suppose that $x \in \varepsilon(\mathbf{J}D\mathbf{U}) = \mathbf{J}D\mathbf{U} \cap X_0$. Then $x \in -\downarrow -\mathbf{I}D\mathbf{U}$, whence $x \notin \downarrow -\mathbf{I}D\mathbf{U}$. It follows that $\uparrow x \cap -\mathbf{I}D\mathbf{U} = \emptyset$. So $\uparrow x \subseteq \mathbf{I}D\mathbf{U} \subseteq \mathbf{D}U = \uparrow \mathbf{C}U$. Therefore, $x \in \uparrow \mathbf{C}U$, and so $\downarrow x \cap \mathbf{C}U \neq \emptyset$. Since $x \in X_0$, then $\downarrow x$ is open. Thus, $\downarrow x \cap U \neq \emptyset$, and as U is an upset of X_0 , it follows that $x \in U$. This proves that $\mathbf{J}D\mathbf{U} \cap X_0 \subseteq U$, whence $\varepsilon(\mathbf{J}D\mathbf{U}) = U$. Consequently, ε is an isomorphism from $\mathcal{RO}(X)$ onto $\text{Up}(X_0)$. By Lemma 3.2, there is an isomorphism $f : \bar{L} \rightarrow \mathcal{RO}(X)$ such that $f \circ \eta = \varphi$. We set $\xi = \varepsilon \circ f$. Then $\xi \circ \eta = \varepsilon \circ f \circ \eta = \varepsilon \circ \varphi = \varphi_0$. Thus, there is an isomorphism $\xi : \bar{L} \rightarrow \text{Up}(X_0)$ such that $\xi \circ \eta = \varphi_0$. \square

Remark 3.8. It follows that if a distributive lattice is principally separated, then its MacNeille completion is distributive (even completely distributive) [5, Sec. 4], thus providing a sufficient condition for the MacNeille completion of a distributive lattice to be distributive. (In fact, each principally separated lattice is distributive. Therefore, that \bar{L} is completely distributive already follows from L being principally separated.)

Let L be a distributive lattice. If L is principally separated, then we saw that \bar{L} is isomorphic to $\text{Up}(Y)$ for some poset Y . But this does not mean that \bar{L} is isomorphic to the lattice of upsets of the Priestley space of L , as the following example shows.




FIGURE 2

Example 3.9. Let L be the negative integers with bottom, and let X be its Priestley space. Then $\langle X, \leq \rangle$ is isomorphic to the positive integers with top, L is complete and principally separated, so $\bar{L} \simeq L$, but $L \not\cong \text{Up}(X)$ (see Fig. 2). Since $\text{Up}(X)$ is isomorphic to L^σ , it follows that $L^\sigma \not\cong \bar{L}$.

Theorem 3.10. *Let L be a distributive lattice and let X be its Priestley space. Then \bar{L} is isomorphic to L^σ iff X_0 is dense in X and $\langle X, \leq \rangle$ is order-isomorphic to $\langle X_0, \leq \rangle$.*

Proof. First suppose that X_0 is dense in X and $\langle X, \leq \rangle$ is order-isomorphic to $\langle X_0, \leq \rangle$. Then $\text{Up}(X_0) \simeq \text{Up}(X)$. By Theorems 3.7 and 3.2, $\bar{L} \simeq \text{Up}(X_0)$ and $L^\sigma \simeq \text{Up}(X)$. Thus, $\bar{L} \simeq L^\sigma$. Now suppose that $\bar{L} \simeq L^\sigma$. Then \bar{L} is completely join-prime generated, and by Theorem 3.7, X_0 is dense in X . Moreover, since $\bar{L} \simeq \text{Up}(X_0)$ and $L^\sigma \simeq \text{Up}(X)$, from $\bar{L} \simeq L^\sigma$ it follows that $\text{Up}(X_0) \simeq \text{Up}(X)$. This by the well-known duality between complete and completely join-prime generated distributive lattices and partially ordered sets implies that $\langle X_0, \leq \rangle$ is order-isomorphic to $\langle X, \leq \rangle$. \square

Corollary 3.11. *Let L be a distributive lattice and let X be its Priestley space. Then the following conditions are equivalent:*

- (1) *The three completions \bar{L} , L^σ , and \widehat{L} of L are isomorphic.*
- (2) *X_0 is dense in X and $\langle X, \leq \rangle$ is order-isomorphic to $\langle X_0, \leq \rangle$.*
- (3) *L is principally separated and the poset $\langle S(L), \geq \rangle$ is order-isomorphic to the poset of prime filters of L .*

Proof. By Lemma 3.2, $L^\sigma \simeq \widehat{L}$. Therefore, the equivalence of (1) and (2) follows from Theorem 3.10. That (2) is equivalent to (3) is a consequence of Theorems 3.5 and 3.4. \square

Example 3.12. To show the utility of Corollary 3.11, we give several concrete examples.

- (1) Let B denote the Boolean lattice of finite and cofinite subsets of ω . It is well-known that the dual space X of B is the one-point compactification of ω , which is homeomorphic to $\omega+1$ in its interval topology (see Fig. 3). Therefore, $X_0 = X_{\text{iso}} = \omega$, and so X_0 is dense in X . Moreover, because both ω and X are countable, and \leq is simply the equality relation, we obtain that $\langle X, \leq \rangle$ is order-isomorphic to $\langle X_0, \leq \rangle$. Consequently, by Corollary 3.11, \bar{B} , B^σ , and \widehat{B} are isomorphic. Note that (\bar{B}, η) , (B^σ, ζ) , and (\widehat{B}, ι) cannot be isomorphic as completions because B is an infinite Boolean algebra (see Theorem 4.8).




FIGURE 3. Example 3.12(1)

(2) Next, consider the Priestley space $\langle X, \tau, \leq \rangle$ shown in Fig. 4, where each (i, j) is




FIGURE 4. Example 3.12(2)

isolated and each (ω, j) is a limit point of X ($i \in \omega$ and $0 \leq j \leq n$). Clearly $X_0 = \{(i, j) \mid i \in \omega \text{ and } 0 \leq j \leq n\}$ and X is the n -point compactification of the discrete space X_0 . Of course, X is a finite disjoint union of the spaces $X^i = \{(n, i) \mid n \in \omega\} \cup \{(\omega, i)\}$, $0 \leq i \leq n$, where each X^i is the one-point compactification of the discrete space $\{(n, i) \mid n \in \omega\}$. We can view this as a generalization of the previous example: we simply replace each point of the previous example by an $(n + 1)$ -point chain. It follows from the definition that X_0 is dense in X , and it is easy to see that $\langle X, \leq \rangle$ is order-isomorphic to $\langle X_0, \leq \rangle$. Let L denote the lattice of clopen upsets of X . Then, by Corollary 3.11, \bar{L} , L^σ , and \hat{L} are isomorphic. By Theorem 3.14, (\bar{L}, η) , (L^σ, ζ) , and (\hat{L}, ι) cannot be isomorphic as completions of L .

(3) Finally, consider the Priestley space $\langle X, \tau, \leq \rangle$ shown in Fig. 5, where each of (i, j) is




FIGURE 5. Example 3.12(3)

isolated, and each of (ω, j) , (i, ω) , and (ω, ω) is a limit point of X ($i, j \in \omega$). Clearly




FIGURE 6

$X_0 = \{(i, j) \mid i, j \in \omega\}$ and X is a countable compactification of the discrete space X_0 . We can think of each $X^i = \{(0, i), (1, i), \dots, (\omega, i)\} \cup \{(i, 0), (i, 1), \dots, (i, \omega)\}$ as the two-point compactification of the discrete space $\{(0, i), (1, i), \dots\} \cup \{(i, 0), (i, 1), \dots\}$, and of X as the one-point compactification of $X - \{(\omega, \omega)\}$ (which is not a discrete space). Therefore, we can view this as a generalization of the previous example: we replace each $(n + 1)$ -point chain by an $(\omega + 1)$ -point chain. It is obvious that X_0 is dense in X . However, $\langle X, \leq \rangle$ is *not* order-isomorphic to $\langle X_0, \leq \rangle$. Let L denote the lattice of clopen upsets of X . Then it follows from Corollary 3.11 that \bar{L} is not isomorphic to neither L^σ nor \widehat{L} .

Based on Example 3.12, it is tempting to conjecture that the MacNeille completion \bar{L} of a distributive lattice L is isomorphic to L^σ and \widehat{L} only if the Priestley space of L does not have any infinite chains. This, however, is not the case as we show in the next example.

Example 3.13. Consider the Priestley space $\langle X, \tau, \leq \rangle$ shown in Fig. 6, where $\{(\omega, n) \mid n \in \omega\} \cup \{(n, \omega) \mid n \in \omega\} \cup \{(\omega, \omega)\}$ are the limit points of X , and the rest are isolated in X . Clearly $X_0 = X - (\{(\omega, n) \mid n \in \omega\} \cup \{(n, \omega) \mid n \in \omega\} \cup \{(\omega, \omega)\})$ and X is a countable compactification of the discrete space X_0 . We can think of each $X^i = \{(0, i), (1, i), \dots, (\omega, i)\} \cup \{(i, 0), (i, 1), \dots, (i, \omega), (i, \omega + i + 1)\}$ as the two-point compactification of the discrete space $X_i - \{(\omega, i), (i, \omega)\}$, and of X as the one point compactification of $X - \{(\omega, \omega)\}$ (which is not a discrete space). Therefore, we can view this as a further generalization of the spaces in Example 3.12. Clearly X has (infinitely many) infinite chains. It is also obvious that X_0 is dense in X . We show that there is an order-isomorphism from




FIGURE 7

X_0 onto X . Define $f: X_0 \rightarrow X$ as follows:

$$f: (\alpha, \beta) \mapsto \begin{cases} (\omega, \beta) & \text{if } \alpha = 0, \beta < \omega, \\ (\omega, \omega) & \text{if } \alpha = 0, \beta = \omega + 1, \\ (\alpha - 1, \beta) & \text{if } \alpha > 0, \beta < \omega, \\ (\alpha - 1, \beta - 1) & \text{if } \alpha > 0, \beta > \omega. \end{cases}$$

The map f is shown in Fig. 7, from which it is easy to see that f is an order-isomorphism. Let L be the lattice of clopen upsets of X . Then, by Corollary 3.11, $\bar{L} \simeq L^\sigma$, in spite of the fact that X has infinite ascending chains. By taking the opposite of X , we obtain an example of a Priestley space X with infinitely descending chains such that X_0 is order-isomorphic to X . Thus, there exist distributive lattices L whose Priestley spaces have both infinitely ascending and descending chains, but still \bar{L} is isomorphic to L^σ (and hence to \widehat{L}).

The final goal of this section is to investigate when (\bar{L}, η) , (L^σ, ζ) , and (\widehat{L}, ι) are isomorphic as completions of L .

Theorem 3.14. *Let L be a distributive lattice. The three completions (\bar{L}, η) , (L^σ, ζ) , and (\widehat{L}, ι) are isomorphic as completions of L iff L is finite.*

Proof. By Lemma 3.2, it is sufficient to show that there is an isomorphism $\alpha: \bar{L} \rightarrow L^\sigma$ such that $\alpha \circ \eta = \zeta$ iff L is finite. Clearly if L is finite, then η and ζ are identities. Therefore, if we let α be the identity, then $\alpha \circ \eta = \zeta$. Conversely, suppose that there exists $\alpha: \bar{L} \rightarrow L^\sigma$ such that $\alpha \circ \eta = \zeta$. Since $\bar{L} \simeq L^\sigma$, then \bar{L} is completely join-prime generated, and by Theorem 3.7, X_0 is dense in X . We show that $X_0 = X$. By Lemma 3.2, $(\bar{L}, \eta) \simeq (\text{Up}(X_0), \varphi_0)$ and $(L^\sigma, \zeta) \simeq (\text{Up}(X), \varphi)$. Therefore, $\alpha \circ \eta = \zeta$ implies there is a lattice isomorphism $\lambda: \text{Up}(X_0) \rightarrow \text{Up}(X)$ such that $\lambda \circ \varphi_0 = \varphi$. Let $x \in X$. Then $\uparrow x \in \text{Up}(X)$. Since λ is onto, there is $U \in \text{Up}(X_0)$ such that $\lambda(U) = \uparrow x$. We have $U = \bigcup \{\uparrow y \cap X_0 \mid y \in U\}$. From $y \in X_0$

it follows that $\uparrow y$ is a clopen upset of X . Therefore, there is $a_y \in L$ such that $\uparrow y = \varphi(a_y)$. Thus,

$$\begin{aligned} U &= \bigcup \{ \uparrow y \cap X_0 \mid y \in U \} \\ &= \bigcup \{ \varphi(a_y) \cap X_0 \mid y \in U \} \\ &= \bigcup \{ \varphi_0(a_y) \mid y \in U \}, \end{aligned}$$

and so

$$\begin{aligned} \uparrow x &= \lambda(U) \\ &= \lambda(\bigcup \{ \varphi_0(a_y) \mid y \in U \}) \\ &= \bigcup \{ \lambda \varphi_0(a_y) \mid y \in U \} \\ &= \bigcup \{ \varphi(a_y) \mid y \in U \}. \end{aligned}$$

Consequently, there is $y \in X_0$ such that $\uparrow x = \varphi(a_y) = \uparrow y$. It follows that $x = y$, and so $x \in X_0$. This implies that $X = X_0$, which together with Lemma 3.6 give us $X_{\text{iso}} = X$. Therefore, X is discrete, hence finite by compactness. Thus, L is finite. \square

We summarize matters below.

Corollary 3.15. *For a distributive lattice L , the three lattices \bar{L} , L^σ , and \widehat{L} are isomorphic iff L is principally separated and the poset $\langle S(L), \geq \rangle$ is order-isomorphic to the poset of prime filters of L . Moreover, the completions (\bar{L}, η) , (L^σ, ζ) , and (\widehat{L}, ι) are isomorphic as completions of L iff L is finite.*

4. HEYTING ALGEBRAS

Recall that a distributive lattice L is a Heyting algebra if there exists a binary operation $\rightarrow: L^2 \rightarrow L$ such that for all $a, b, c \in L$ we have:

$$a \wedge c \leq b \text{ iff } c \leq a \rightarrow b.$$

The duality theory for Heyting algebras was developed by Esakia [7]. As we will see shortly, the MacNeille and canonical completions of a Heyting algebra have the same dual characterization as for distributive lattices. Therefore, our results about the isomorphism of \bar{L} and L^σ apply unchanged for Heyting algebras. Some simplifications are possible though, because for a Heyting algebra A we have $S(A) = J^\infty(A)$. On the other hand, the dual characterization of the profinite completion of a Heyting algebra is different from that of a distributive lattice. In this section we will adjust the results of the previous section appropriately to obtain necessary and sufficient conditions for the MacNeille, canonical, and profinite completions of a Heyting algebra to be isomorphic. As a corollary, we also obtain necessary and sufficient conditions for the three completions of a Boolean algebra to be isomorphic.

Definition 4.1. [7] *A triple $\langle X, \tau, \leq \rangle$ is an Esakia space if $\langle X, \tau, \leq \rangle$ is a Priestley space and $\downarrow U$ is open for each open subset U of X .*

The same way distributive lattices are represented as clopen upsets of Priestley spaces, Heyting algebras are represented as clopen upsets of Esakia spaces.

We recall that there is a 1-1 correspondence between congruences and filters of a Heyting algebra A (see, e.g., [19, Section I.13]), and that it is obtained by associating with each congruence θ of A , the filter $F_\theta = 1/\theta = \{a \in A \mid a\theta 1\}$, and with each filter F of A , the

congruence $\theta_F = \{(a, b) \in A^2 \mid (a \rightarrow b) \wedge (b \rightarrow a) \in F\}$. We say that a filter F of A is of *finite index* if the quotient algebra A/θ_F is finite.

Let A be a Heyting algebra and let X be its Esakia space. We recall that filters of A correspond to closed upsets of X (see, e.g., [7]), and that this correspondence is obtained by associating with each filter F of A , the closed upset $C_F = \bigcap \{\varphi(a) \mid a \in F\}$ of X , and with each closed upset C of X , the filter $F_C = \{a \in A \mid C \subseteq \varphi(a)\}$ of A . Now, if F is a principal filter $\uparrow a$ of A , then $C_F = \varphi(a)$ is a clopen upset of X , and if C is a clopen upset of X , then $C = \varphi(a)$ for some $a \in A$, and so F_C is the principal filter $\uparrow a$ of A . Thus, principal filters of A correspond to clopen upsets of X . Also, if F is of finite index, then C_F is a finite upset of X , and if C is a finite upset of X , then F_C is of finite index. Thus, filters of A of finite index correspond to finite upsets of X .

As we pointed out above, in a Heyting algebra A we have $S(A) = J^\infty(A)$, and so for Heyting algebras principally separated simply means completely join-prime generated. This follows from [2, Thm. 2.7] and the dual characterization of splitting pairs of a distributive lattice given in Theorem 3.4. We also give a simple algebraic proof. Let $a \in J^\infty(A)$. If $\downarrow a - \{a\}$ does not have a greatest element, then $a = \bigvee(\downarrow a - \{a\})$, which contradicts to $a \in J^\infty(A)$. Let b be the greatest element of $\downarrow a - \{a\}$. Then $(a, a \rightarrow b)$ splits A . Indeed, for $c \in A$, if $a \not\leq c$, then $a \wedge c \leq b$, so $c \leq a \rightarrow b$, and so $a \in S(A)$.

Let X be an Esakia space. Since for $x \in X$ we have $x \in X_{\text{iso}}$ implies $\downarrow x$ is clopen, we obtain that order-isolated points of X are exactly those isolated points x of X for which $\uparrow x$ is (cl)open. We define

$$X_{\text{fin}} = \{x \in X \mid \uparrow x \text{ is finite}\}.$$

In other words, X_{fin} is the union of finite upsets of X . Let $\varphi_{\text{fin}}: A \rightarrow \text{Up}(X_{\text{fin}})$ be given by $\varphi_{\text{fin}}(a) = \varphi(a) \cap X_{\text{fin}}$. Then φ_{fin} is a lattice homomorphism, and φ_{fin} is an embedding iff X_{fin} is dense in X [2, Theorem 3.1 and Proposition 3.2].

Lemma 4.2. *Let A be a Heyting algebra and let X be its Esakia space. Then:*

- (1) *There is an isomorphism $f: \bar{A} \rightarrow \mathcal{RO}(X)$ such that $f \circ \eta = \varphi$. Therefore, $(\bar{A}, \eta) \simeq (\mathcal{RO}(X), \varphi)$. Moreover, $(\bar{A}, \eta) \simeq (\text{Up}(X_0), \varphi_0)$ iff A is completely join-prime generated.*
- (2) *There is an isomorphism $g: A^\sigma \rightarrow \text{Up}(X)$ such that $g \circ \zeta = \varphi$. Therefore, $(A^\sigma, \zeta) \simeq (\text{Up}(X), \varphi)$.*
- (3) *There is an isomorphism $h: \hat{A} \rightarrow \text{Up}(X_{\text{fin}})$ such that $h \circ \iota = \varphi_{\text{fin}}$. Therefore, if $\iota: A \rightarrow \hat{A}$ is an embedding, then $(\hat{A}, \iota) \simeq (\text{Up}(X_{\text{fin}}), \varphi_{\text{fin}})$.*

Proof. Since A is principally separated iff A is completely join-prime generated, (1) follows from Lemma 3.2 and Theorem 3.7. Obviously (2) is a consequence of Lemma 3.2. To see (3), it was shown in [3, Theorem 4.7] that $\hat{A} \simeq \text{Up}(X_{\text{fin}})$. We recall that the isomorphism $h: \hat{A} \rightarrow \text{Up}(X_{\text{fin}})$ is explicitly defined as follows. Let Ψ_A denote the set of filters of A of finite index. For $F \in \Psi_A$, let C_F be the finite upset of X corresponding to F . For $\alpha \in \hat{A}$ and $F \in \Psi_A$, we have $\alpha(F) = a/\theta_F$ for some $a \in A$. Then $\varphi(a) \cap C_F$ is an upset of C_F . If there is another $b \in L$ such that $\alpha(F) = b/\theta_F$, then $\varphi(a) \cap C_F = \varphi(b) \cap C_F$. Thus, $\varphi(a) \cap C_F$ is independent of $a \in L$, and we denote it by $\varphi(\alpha(F))$. Let $U_\alpha = \bigcup \{\varphi(\alpha(F)) \mid F \in \Psi_A\}$. Since each C_F is a finite upset of X and $\varphi(\alpha(F))$ is an upset of C_F , it follows that U_α is an upset of X_{fin} . We define $h: \hat{A} \rightarrow \text{Up}(X_{\text{fin}})$ by $h(\alpha) = U_\alpha$. Then h is the desired isomorphism. We show that $h \circ \iota = \varphi_{\text{fin}}$. For $a \in A$ we have $h(\iota(a)) = h(\alpha_a) = \bigcup \{\varphi(a) \cap C_F \mid F \in \Psi_A\} =$

$\varphi(a) \cap \bigcup \{C_F \mid F \in \Psi_A\} = \varphi(a) \cap X_{\text{fin}} = \varphi_{\text{fin}}(a)$. Here we use that $X_{\text{fin}} = \bigcup \{C_F \mid F \in \Psi_A\}$, which is true because X_{fin} is the union of finite upsets of X and each finite upset of X has the form C_F for some $F \in \Psi_A$. \square

From Theorem 3.10 and Lemma 4.2 we immediately obtain:

Theorem 4.3. *Let A be a Heyting algebra and let X be its Esakia space. Then the following conditions are equivalent:*

- (1) $\overline{A} \simeq A^\sigma$.
- (2) X_0 is dense in X and $\langle X_0, \leq \rangle$ is order-isomorphic to $\langle X, \leq \rangle$.
- (3) A is completely join-prime generated and $\langle J^\infty(A), \geq \rangle$ is order-isomorphic to the poset of prime filters of A .

Since the profinite completion of a Heyting algebra behaves differently from the profinite completion of a distributive lattice, the comparison of the MacNeille and canonical completions of a Heyting algebra to its profinite completion is slightly different.

Theorem 4.4. *Let A be a Heyting algebra and let X be its Esakia space. Then:*

- (1) *The following conditions are equivalent:*
 - (a) $\overline{A} \simeq \widehat{A}$.
 - (b) X_0 is dense in X and $\langle X_0, \leq \rangle$ is order-isomorphic to $\langle X_{\text{fin}}, \leq \rangle$.
 - (c) A is completely join-prime generated and $\langle J^\infty(A), \geq \rangle$ is order-isomorphic to the poset of prime filters of A of finite index.
- (2) *The following conditions are equivalent:*
 - (a) $A^\sigma \simeq \widehat{A}$.
 - (b) $(A^\sigma, \zeta) \simeq (\widehat{A}, \iota)$.
 - (c) $\langle X_{\text{fin}}, \leq \rangle$ is order-isomorphic to $\langle X, \leq \rangle$.
 - (d) $X_{\text{fin}} = X$.

Proof. The proof of (1) is analogous to that of Theorem 3.10. For (2), note that the equivalences (a) \Leftrightarrow (c) \Leftrightarrow (d) follow from [3, Theorem 4.10]. Furthermore, it is clear that (b) implies (a). Finally, (d) implies (b) because from $X_{\text{fin}} = X$ it follows that $\text{Up}(X_{\text{fin}}) = \text{Up}(X)$ and the identity $\text{id}_{\text{Up}(X)}: \text{Up}(X) \rightarrow \text{Up}(X)$ is always an isomorphism of completions. \square

Putting Theorems 4.3 and 4.4 together, we obtain:

Corollary 4.5. *Let A be a Heyting algebra and let X be its Esakia space. Then the following conditions are equivalent:*

- (1) $\overline{A} \simeq A^\sigma \simeq \widehat{A}$.
- (2) X_0 is dense in X and $\langle X_0, \leq \rangle$ is order-isomorphic to $\langle X_{\text{fin}}, \leq \rangle = \langle X, \leq \rangle$.
- (3) A is completely join-prime generated, $\langle J^\infty(A), \geq \rangle$ is isomorphic to the poset of prime filters of A , and every prime filter of A is of finite index.

We mention some of the consequences of these results below. We say that a Heyting algebra A is *finitely approximable*, or *residually finite*, if $a \not\leq b$ implies there is a filter F of A of finite index such that $a \in F$ and $b \notin F$.

Corollary 4.6. *Let A be a Heyting algebra and let X be its Esakia space.*

- (1) *If the variety generated by A is finitely generated, then $\overline{A} \simeq A^\sigma \simeq \widehat{A}$ iff X_0 is dense in X and $\langle X_0, \leq \rangle$ is order-isomorphic to $\langle X, \leq \rangle$.*




FIGURE 8

- (2) If X has infinite chains, then A^σ is not isomorphic to \widehat{A} . Nevertheless, there are Heyting algebras whose Esakia spaces have infinite chains but \overline{A} is still isomorphic to A^σ ; also, there are Heyting algebras whose Esakia spaces have infinite chains but \overline{A} is still isomorphic to \widehat{A} .
- (3) If A is a finitely generated and finitely approximable Heyting algebra, then \overline{A} is isomorphic to \widehat{A} .

Proof. (1) follows from Corollary 4.5 and the fact that $X_{\text{fin}} = X$ whenever A generates a finitely generated variety of Heyting algebras (see, e.g., [3, Theorem 5.1]). For (2), observe that if X has infinite chains, then $X_{\text{fin}} \neq X$, and apply Theorem 4.4.2. For an example of a Heyting algebra A whose Esakia space has infinite chains but \overline{A} is still isomorphic to \widehat{A} , observe that the Priestley space constructed in Example 3.13 is in fact an Esakia space. Therefore, the corresponding Heyting algebra is our desired example. Finally, for an example of a Heyting algebra A whose Esakia space has infinite chains but \overline{A} is still isomorphic to \widehat{A} take the Heyting algebra A of positive integers with top. The Esakia space X of A is order-isomorphic to the negative integers with bottom (see Fig. 8). So $X_0 = X_{\text{fin}}$, and so $\overline{A} \simeq \widehat{A} \simeq A$. For (3) observe that if A is a finitely generated and finitely approximable Heyting algebra, then it follows from [4, Section 3.1] and [2, Theorem 3.1] that $X_0 = X_{\text{iso}} = X_{\text{fin}}$ and that X_0 is dense in X , thus the result follows from Theorem 4.4. \square

Corollary 4.7. *For a Boolean algebra A , the following conditions are equivalent:*

- (1) $\overline{A} \simeq A^\sigma \simeq \widehat{A}$.
- (2) A is atomic and the set of atoms of A has the same cardinality as the set of ultrafilters of A .
- (3) A is atomic and the cardinality of the set of free ultrafilters of A is less than or equal to the cardinality of the set of atoms of A .

Proof. Suppose A is a Boolean algebra and X is its Stone space. Then $X_0 = X_{\text{iso}}$, there is a bijection between atoms of A and X_{iso} , and A is atomic iff X_{iso} is dense in X . Moreover, an order-isomorphism between X_0 and X is simply a bijection. Now since the variety of Boolean algebras is a finitely generated variety of Heyting algebras, the result follows from Corollary 4.6.1. \square

The rest of this section we will be concerned with necessary and sufficient conditions for the MacNeille, canonical, and profinite completions of a Heyting algebra A to be isomorphic as completions of A . First off, as a reformulation of Theorem 3.14 we obtain:

Theorem 4.8. *For a Heyting algebra A , there is an isomorphism of completions $\alpha: (\overline{A}, \eta) \rightarrow (A^\sigma, \zeta)$ iff A is finite. In particular, if A is a Boolean algebra, then $(\overline{A}, \eta) \simeq (A^\sigma, \zeta)$ iff A is finite.*

Now we obtain a necessary and sufficient condition for an isomorphism $\beta: \overline{A} \rightarrow \widehat{A}$ to commute with η and ι .

Lemma 4.9. *Let $\langle X, \tau \rangle$ be a topological space with $Y \subseteq X$ dense in X . If $Z \subseteq X$ is open and $Z \cap Y$ is closed in X , then $Z \subseteq Y$.*

Proof. Let τ_Z denote the subspace topology on Z and let \mathbf{C}_Z denote the closure operator in $\langle Z, \tau_Z \rangle$. Since Z is open in $\langle X, \tau \rangle$, we have $Z \cap Y$ is dense in $\langle Z, \tau_Z \rangle$. Therefore, $\mathbf{C}_Z(Z \cap Y) = Z$. Because $Z \cap Y$ is closed in $\langle X, \tau \rangle$, it is also closed in $\langle Z, \tau_Z \rangle$. Thus, $Z \cap Y = \mathbf{C}_Z(Z \cap Y) = Z$. It follows that $Z \subseteq Y$. \square

Lemma 4.10. *Let A be a Heyting algebra and let X be its Esakia space. If $\overline{A} \simeq \widehat{A}$, then $X_0 = X_{\text{iso}} \subseteq X_{\text{fin}}$. Consequently, X_{fin} is dense and A is finitely approximable.*

Proof. Since $\overline{A} \simeq \widehat{A}$, it follows from Theorem 4.4.1 that X_0 is dense in X , and that $\langle X_0, \leq \rangle$ is order-isomorphic to $\langle X_{\text{fin}}, \leq \rangle$. As X_0 is dense in X , by Lemma 3.6, $X_0 = X_{\text{iso}}$. From $\langle X_0, \leq \rangle \simeq \langle X_{\text{fin}}, \leq \rangle$ it follows that $\uparrow x \cap X_0$ is finite for each $x \in X_0$. Since X is Hausdorff, $\uparrow x \cap X_0$ is closed. Moreover, because $x \in X_0$, $\uparrow x$ is open by definition; so by Lemma 4.9, $\uparrow x \subseteq X_0$. Therefore, $\uparrow x = \uparrow x \cap X_0$ is finite, whence $x \in X_{\text{fin}}$. It follows that $X_0 \subseteq X_{\text{fin}}$, so X_{fin} is dense, and so, by [2, Theorem 3.1], A is finitely approximable. \square

Note that $\overline{A} \simeq \widehat{A}$ does not imply that X_0 is equal to X_{fin} as the following example shows.

Example 4.11. Let B be the Boolean algebra of finite and cofinite subsets of ω and let X be its Stone space. Then X is homeomorphic to the one-point compactification of ω (see Fig. 3), $X_0 = X_{\text{iso}} = \omega$ and it is a proper subset of $X_{\text{fin}} = X$. On the other hand, both \overline{B} and \widehat{B} are isomorphic to the powerset of ω . Since $B^\sigma \simeq \mathcal{P}(X) \simeq \mathcal{P}(\omega)$, because both ω and X are countable, it follows that $\overline{B} \simeq \widehat{B} \simeq B^\sigma$.

Lemma 4.12. *Let A be a Heyting algebra and let X be its Esakia space. Then the following conditions are equivalent:*

- (1) $X_{\text{fin}} \subseteq X_0$.
- (2) *Each filter of A of finite index is principal.*

Proof. (1) \Rightarrow (2): Let F be a filter of A and let C_F be the corresponding closed upset of X . If F is of finite index, then A/θ_F is finite, and so C_F is also finite. Therefore, by (1), C_F is a finite subset of X_0 , so is clopen. Thus, F is a principal filter.

(2) \Rightarrow (1): Let $x \in X_{\text{fin}}$. Then $\uparrow x$ is a finite closed upset of X . Therefore, the corresponding filter of A is of finite index, hence principal by (2). It follows that $\uparrow x$ is a finite clopen upset of X . Thus, x is an isolated point of X and $\uparrow x$ is clopen, so $x \in X_0$. \square

As follows from [4, Chapter 3], if A is the free n -generated Heyting algebra and X is its Esakia space, then $X_{\text{fin}} \subseteq X_0$, so each filter of A of finite index is principal.

Theorem 4.13. *Let A be a Heyting algebra and let X be its Esakia space. Then the following conditions are equivalent:*

- (1) *There is an isomorphism of completions $\gamma: (\overline{A}, \eta) \rightarrow (\widehat{A}, \iota)$.*
- (2) *X_0 is dense and $X_0 = X_{\text{fin}}$.*
- (3) *A is finitely approximable and each filter of A of finite index is principal.*

Proof. (1) \Rightarrow (2): Suppose that there is an isomorphism $\gamma: \overline{A} \rightarrow \widehat{A}$ such that $\gamma \circ \eta = \iota$. Then $\overline{A} \simeq \widehat{A}$, so by Theorem 4.4 and Lemma 4.10, X_0 is dense and $X_0 \subseteq X_{\text{fin}}$. It is left to be shown that $X_{\text{fin}} \subseteq X_0$. By Lemma 4.2, $(\overline{A}, \eta) \simeq (\text{Up}(X_0), \varphi_0)$ and $(\widehat{A}, \iota) \simeq (\text{Up}(X_{\text{fin}}), \varphi_{\text{fin}})$. Therefore, $\gamma \circ \eta = \iota$ implies there is a lattice isomorphism $\lambda: \text{Up}(X_0) \rightarrow \text{Up}(X_{\text{fin}})$ such that $\lambda \circ \varphi_0 = \varphi_{\text{fin}}$. Let $x \in X_{\text{fin}}$. Then $\uparrow x \in \text{Up}(X_{\text{fin}})$. Since λ is onto, there is $U \in \text{Up}(X_0)$ such that $\lambda(U) = \uparrow x$. We have $U = \bigcup \{\uparrow y \cap X_0 \mid y \in U\}$. From $y \in X_0$ it follows that $\uparrow y$ is a clopen upset of X . Therefore, there is $a_y \in A$ such that $\uparrow y = \varphi(a_y)$. Thus,

$$\begin{aligned} U &= \bigcup \{\uparrow y \cap X_0 \mid y \in U\} \\ &= \bigcup \{\varphi(a_y) \cap X_0 \mid y \in U\} \\ &= \bigcup \{\varphi_0(a_y) \mid y \in U\}, \end{aligned}$$

and so

$$\begin{aligned} \uparrow x &= \lambda(U) \\ &= \lambda(\bigcup \{\varphi_0(a_y) \mid y \in U\}) \\ &= \bigcup \{\lambda \varphi_0(a_y) \mid y \in U\} \\ &= \bigcup \{\varphi_{\text{fin}}(a_y) \mid y \in U\}. \end{aligned}$$

Consequently, there is $y \in X_0$ such that $\uparrow x = \varphi_{\text{fin}}(a_y) = \varphi(a_y) \cap X_{\text{fin}} = \uparrow y \cap X_{\text{fin}} = \uparrow y$ since $y \in X_0 \subseteq X_{\text{fin}}$. It follows that $x = y$, so $x \in X_0$, and so $X_0 = X_{\text{fin}}$.

(2) \Rightarrow (1): If X_0 is dense, then by Theorem 3.7, there is an isomorphism $f: \overline{A} \rightarrow \text{Up}(X_0)$ such that $f \circ \eta = \varphi_0$. From $X_0 = X_{\text{fin}}$ it follows that X_{fin} is dense, $\text{Up}(X_0) = \text{Up}(X_{\text{fin}})$, and $\varphi_0 = \varphi_{\text{fin}}$. By Lemma 4.2, there is an isomorphism $h: \widehat{A} \rightarrow \text{Up}(X_{\text{fin}})$ such that $h \circ \iota = \varphi_{\text{fin}}$. Let $\gamma = h^{-1} \circ f$. Then $\gamma: \overline{A} \rightarrow \widehat{A}$ is an isomorphism such that $\gamma \circ \eta = \iota$.

(2) \Rightarrow (3): If X_0 is dense and $X_0 = X_{\text{fin}}$, then X_{fin} is also dense, hence A is finitely approximable by [2, Theorem 3.1]. Moreover, $X_{\text{fin}} \subseteq X_0$ implies, by Lemma 4.12, that each filter of A of finite index is principal.

(3) \Rightarrow (2): If A is finitely approximable, then X_{fin} is dense by [2, Theorem 3.1]. Therefore, $X_0 \subseteq X_{\text{iso}} \subseteq X_{\text{fin}}$. Moreover, since each filter of A of finite index is principal, by Lemma 4.12, $X_{\text{fin}} \subseteq X_0$. Thus, $X_0 = X_{\text{fin}}$ and X_0 is dense. \square

Among the examples of Heyting algebras that satisfy conditions of Theorem 4.13 are finitely generated finitely approximable Heyting algebras (cf. Corollary 4.6.3). Thus, for finitely generated finitely approximable Heyting algebras A , the strengthening of Corollary 4.6.3 to the isomorphism of \overline{A} and \widehat{A} that commutes with η and ι is an immediate consequence of Theorem 4.13. Another immediate consequence of Theorems 4.13 and 4.8 is the following:

Corollary 4.14. *For a Heyting algebra A , $(\overline{A}, \eta) \simeq (A^\sigma, \zeta) \simeq (\widehat{A}, \iota)$ iff A is finite. In particular, if A is a Boolean algebra, then $(\overline{A}, \eta) \simeq (A^\sigma, \zeta) \simeq (\widehat{A}, \iota)$ iff A is finite.*

We conclude by summarizing the main theorems of this section.

Corollary 4.15. *Let A be a Heyting algebra.*

- (1) (a) $\bar{A} \simeq A^\sigma$ iff A is completely join-prime generated and $\langle J^\infty(A), \geq \rangle$ is order-isomorphic to the poset of prime filters of A .
 (b) $(\bar{A}, \eta) \simeq (A^\sigma, \zeta)$ iff A is finite.
- (2) (a) $\bar{A} \simeq \hat{A}$ iff A is completely join-prime generated and $\langle J^\infty(A), \geq \rangle$ is order-isomorphic to the poset of prime filters of A of finite index.
 (b) $(\bar{A}, \eta) \simeq (\hat{A}, \iota)$ iff A is finitely approximable and each filter of A of finite index is principal.
- (3) $A^\sigma \simeq \hat{A}$ iff $(A^\sigma, \zeta) \simeq (\hat{A}, \iota)$ iff each prime filter of A is of finite index.
- (4) (a) $\bar{A} \simeq A^\sigma \simeq \hat{A}$ iff A is completely join-prime generated, $\langle J^\infty(A), \geq \rangle$ is isomorphic to the poset of prime filters of A , and each prime filter of A is of finite index.
 (b) $(\bar{A}, \eta) \simeq (A^\sigma, \zeta) \simeq (\hat{A}, \iota)$ iff A is finite.

5. FURTHER DIRECTIONS

The results of Section 4 can easily be modified to apply to modal algebras in light of the fact that the Jónsson-Tarski duality for modal algebras is essentially the Esakia duality formulated for modal algebras. The modal algebra case of Theorem 4.13 and Corollary 4.6.3 will be described in detail in [20].

More generally, using the duality theory developed by Goldblatt [11], we would expect that our main results generalize to the case of distributive lattices with operators. In fact, such results as those in [8] and [12] indicate that distributivity of the underlying lattice may not be essential at all.

ACKNOWLEDGMENT

We are greatly indebted to the referee for many valuable suggestions. More specifically, the notion of a principally separated lattice (Definition 2.4.1) and the fact that there exist (incomplete) completely join-prime generated lattices, which are not principally separated (Example 2.3) were suggested by the referee. In addition, his/her comments resulted in the reformulations of Lemmas 3.2 and 4.2 and Theorems 3.4, 3.5, and 3.7. He/she also pointed out some inaccuracies in the original proofs of Theorems 3.7, 3.14, and 4.13.

REFERENCES

- [1] B. Banaschewski and G. Bruns. Categorical characterization of the MacNeille completion. *Archiv der Mathematik*, 18:369–377, 1967.
- [2] G. Bezhanishvili and N. Bezhanishvili. Profinite Heyting algebras. *Order*, 2008. To appear, also available at <http://www.cs.le.ac.uk/people/nb118/Publications/Bezh-Bezh-final.pdf>.
- [3] G. Bezhanishvili, M. Gehrke, R. Mines, and P.J. Morandi. Profinite completions and canonical extensions of Heyting algebras. *Order*, 23:143–161, 2006.
- [4] N. Bezhanishvili. *Lattices of Intermediate and Cylindric Modal Logics*. PhD thesis, University of Amsterdam, 2006.
- [5] Marcel Ern e. Bigeneration in complete lattices and principal separation in ordered sets. *Order*, 8(2):197–221, 1991.
- [6] Marcel Ern e. The Dedekind-MacNeille completion as a reflector. *Order*, 8(2):159–173, 1991.
- [7] L. L. Esakia. Topological Kripke models. *Soviet Math. Dokl.*, 15:147–151, 1974.
- [8] M. Gehrke and J. Harding. Bounded lattice expansions. *Journal of Algebra*, 239:345–371, 2001.
- [9] M. Gehrke, J. Harding, and Y. Venema. MacNeille completions and canonical extensions. *Transactions of the American Mathematical Society*, 358:573–590, 2006.

- [10] M. Gehrke and B. Jónsson. Bounded distributive lattices with operators. *Mathematica Japonica*, 2:207–215, 1994.
- [11] R. Goldblatt. Varieties of complex algebras. *Annals of Pure and Applied Logic*, 44:173–242, 1989.
- [12] J. Harding. On profinite completions and canonical extensions. *Algebra Universalis*, 55:293–296, 2006.
- [13] J. Harding and G. Bezhanishvili. MacNeille completions of Heyting algebras. *Houston Journal of Mathematics*, 30:937–952, 2004.
- [14] B. Jónsson and A. Tarski. Boolean algebras with operators. Part I. *American Journal of Mathematics*, 73:891–939, 1951.
- [15] Ralph McKenzie. Equational bases and nonmodular lattice varieties. *Trans. Amer. Math. Soc.*, 174:1–43, 1972.
- [16] H. Ono. Algebraic semantics for predicate logics and their completeness. In *Logic at work*, volume 24 of *Stud. Fuzziness Soft Comput.*, pages 637–650. Physica, Heidelberg, 1999.
- [17] H. Ono. Completions of algebras and completeness of modal and substructural logics. In *Advances in modal logic. Vol. 4*, pages 335–353. King’s Coll. Publ., London, 2003.
- [18] H.A. Priestley. Representation of distributive lattices by means of ordered Stone spaces. *Bulletin of the London Mathematical Society*, 2:186–190, 1970.
- [19] H. Rasiowa and R. Sikorski. *The Mathematics of Metamathematics*. Monografie Matematyczne, Tom 41. Państwowe Wydawnictwo Naukowe, Warsaw, 1963.
- [20] J. Vosmaer. MacNeille completion and profinite completion can coincide on finitely generated modal algebras. To appear in *Algebra Universalis*, 2008.
- [21] Phillip M. Whitman. Splittings of a lattice. *Amer. J. Math.*, 65:179–196, 1943.

Guram Bezhanishvili
Department of Mathematical Sciences
New Mexico State University
Las Cruces NM 88003-8001, USA
Email: gbezhani@nmsu.edu

Jacob Vosmaer
Institute for Logic, Language and Computation
University of Amsterdam
Plantage Muidergracht 24, 1018 TV Amsterdam
The Netherlands
J.Vosmaer@uva.nl