

MACNEILLE COMPLETION AND PROFINITE COMPLETION CAN COINCIDE ON FINITELY GENERATED MODAL ALGEBRAS

JACOB VOSMAER

ABSTRACT. Following Bezhanishvili & Vosmaer, we confirm a conjecture of Yde Venema by piecing together results from various authors. Specifically, we show that if \mathbb{A} is a residually finite, finitely generated modal algebra such that $\text{HSP}(\mathbb{A})$ has equationally definable principal congruences, then the profinite completion of \mathbb{A} is isomorphic to its MacNeille completion, and \diamond is smooth. Specific examples of such modal algebras are the free **K4**-algebra and the free **PDL**-algebra.

1. INTRODUCTION

In this paper we compare two mathematical constructions applied to modal algebras. The first is the MacNeille completion, which is an order-theoretic generalization of the construction of the reals from the rationals using Dedekind cuts [12]. It has been applied in logic to e.g. prove the completeness of predicate calculi [14]. The second is the profinite completion, which is a universal algebraic construction, transforming an algebra into a topological algebra endowed with a Stone (compact, Hausdorff, zero-dimensional) topology. This construction stems from Galois theory [15], but has more recently also been connected with lattice completions [4, 10, 19, 3].

This paper is a companion piece to [3]. In that paper, parallel versions of our Theorems 2 and 4 arise in a study of the connections between different completions of Heyting algebras, using Esakia duality. In light of the topological character of the profinite completion, in the present paper we will present topological algebra proofs instead. This establishes a strong connection with the body of work on canonicity [8] and MacNeille canonicity [17]. Another advantage of our present perspective is that we can show how the two main Theorems pivot around an interaction between principal lattice filters and principal algebra congruences, which are in a 1-1 correspondence for Heyting algebras, but not for modal algebras. Finally, we will briefly mention some of the connections of our results to modal logic.

The author would like to thank Yde Venema, who suggested that Theorem 4 might be true. Additionally, the author is grateful to the Editor and the Referee for their criticisms and suggestions.

Date: April 28, 2008.

2000 Mathematics Subject Classification. Primary 06E25; Secondary 06B23, 03B45, 22A30.

Key words and phrases. modal algebra, MacNeille completion, profinite completion.

This research was supported by VICI grant 639.073.501 of the Netherlands Organization for Scientific Research (NWO).

2. COMPLETIONS AND TOPOLOGIES

Let $\mathbb{B} = \langle B; \wedge, \vee, \neg, 0, 1 \rangle$ be a Boolean algebra. Given $b \in B$ we write $b\downarrow = \{a \in B \mid a \leq b\}$ ($b\uparrow$ is defined dually). We say $S \subseteq B$ is *join-dense* in \mathbb{B} iff for every $a \in B$, $a = \bigvee(a\downarrow \cap S)$ (*meet-density* is defined dually). A *completion* of a lattice \mathbb{B} is a pair (m, \mathbb{C}) , where $m: \mathbb{B} \hookrightarrow \mathbb{C}$ is a lattice embedding into a complete lattice \mathbb{C} . Completions (m, \mathbb{C}) and (k, \mathbb{D}) of \mathbb{B} are isomorphic if $gm = k$ for some lattice isomorphism $g: \mathbb{C} \rightarrow \mathbb{D}$. If (m, \mathbb{C}) is a completion of \mathbb{B} , let $\rho_{\mathbb{B}}$ be the topology on \mathbb{C} generated by basis $\{[m(a), m(b)] \mid a, b \in B\}$ (where $[x, y] = \{z \in \mathbb{C} \mid x \leq z \leq y\}$). By $\gamma_{\mathbb{B}}^{\downarrow}, \gamma_{\mathbb{B}}^{\uparrow}$ and $\gamma_{\mathbb{B}}$ we denote the Scott topology, the dual Scott topology, and the biScott topology on \mathbb{B} respectively. Let $\text{At } \mathbb{B}$ be the (possibly empty) set of atoms of \mathbb{B} , and let $\text{At}_{\omega} \mathbb{B}$ be the set of all finite joins of atoms of \mathbb{B} . Then $\iota_{\mathbb{B}}$ is the topology generated by the basis $\{[a, \neg b] \mid a, b \in \text{At}_{\omega} \mathbb{B}\}$. By [8, Section 2], $\iota_{\mathbb{B}} = \gamma_{\mathbb{B}}$ if \mathbb{B} is complete and atomic.

The *MacNeille completion* [2] of a Boolean algebra \mathbb{B} is defined up to isomorphism as a completion (m, \mathbb{C}) such that $m[B]$ is join-dense in \mathbb{C} (by [5, Theorem V-27] \mathbb{C} is then also a Boolean algebra). We denote the MacNeille completion of \mathbb{B} by $\bar{\mathbb{B}}$. Alternatively [17, Theorem 4.5], $\bar{\mathbb{B}}$ can be characterized up to isomorphism as a completion (m, \mathbb{C}) of \mathbb{B} such that $\langle \mathbb{C}, \rho_{\mathbb{B}} \rangle$ is Hausdorff. If $f: \mathbb{B} \rightarrow \mathbb{C}$ is an order-preserving map between Boolean algebras, then $f^{\circ}: \bar{\mathbb{B}} \rightarrow \bar{\mathbb{C}}$, defined by $f^{\circ}: x \mapsto \bigvee \{f(a) \mid m_{\mathbb{B}}(a) \leq x\}$, is the *lower extension* of f . The *upper extension* f^{\bullet} is defined dually. Alternatively [17, Section 5], f° is the (pointwise) largest $(\rho_{\mathbb{B}}, \gamma_{\bar{\mathbb{C}}}^{\downarrow})$ -continuous extension of f , and f^{\bullet} is the smallest $(\rho_{\mathbb{B}}, \gamma_{\bar{\mathbb{C}}}^{\uparrow})$ -continuous extension of f . We say f is *smooth* if $f^{\circ} = f^{\bullet}$.

Given a modal algebra $\mathbb{A} = \langle A; \diamond \rangle$, let $\Phi_{\mathbb{A}} := \{\theta \in \text{Con } \mathbb{A} \mid \mathbb{A}/\theta \text{ is finite}\}$. We say \mathbb{A} is *residually finite* if for all $a, b \in \mathbb{A}$ with $a \neq b$, there exists $\theta \in \Phi_{\mathbb{A}}$ such that $a/\theta \neq b/\theta$. The inverse system $\langle \{\mathbb{A}/\theta\}_{\theta \in \Phi_{\mathbb{A}}}, f_{\theta\psi} \rangle$, where $f_{\theta\psi}: \mathbb{A}/\theta \rightarrow \mathbb{A}/\psi$ (for all $\theta, \psi \in \Phi_{\mathbb{A}}$ such that $\theta \subseteq \psi$) is defined by $f_{\theta\psi}: a/\theta \mapsto a/\psi$, has a projective limit

$$\hat{\mathbb{A}} = \{\alpha \in \prod_{\theta \in \Phi_{\mathbb{A}}} \mathbb{A}/\theta \mid \forall \theta, \psi \in \Phi_{\mathbb{A}} \text{ with } \theta \subseteq \psi, \text{ if } \alpha(\theta) = a/\theta \text{ then } \alpha(\psi) = a/\psi\}.$$

The map $\mu: \mathbb{A} \rightarrow \hat{\mathbb{A}}$, defined by $\mu: a \mapsto (a/\theta)_{\theta \in \Phi_{\mathbb{A}}}$, is a modal algebra homomorphism which is injective iff \mathbb{A} is residually finite. We call $\hat{\mathbb{A}}$ the *profinite completion* of \mathbb{A} [15]. Since $\hat{\mathbb{A}}$ is a complete lattice [10], it follows that $(\mu, \hat{\mathbb{A}})$ is a completion of \mathbb{A} iff \mathbb{A} is residually finite. If we define the discrete topology on each \mathbb{A}/θ , $\hat{\mathbb{A}}$ inherits a topology $\tau_{\hat{\mathbb{A}}}$ as a closed subspace of the product $\prod_{\theta \in \Phi_{\mathbb{A}}} \mathbb{A}/\theta$. Now $\langle \hat{\mathbb{A}}, \tau_{\hat{\mathbb{A}}} \rangle$ is a Stone space [4, Section 2], and in particular $\hat{\diamond}: \hat{\mathbb{A}} \rightarrow \hat{\mathbb{A}}$ is $(\tau_{\hat{\mathbb{A}}}, \tau_{\hat{\mathbb{A}}})$ -continuous [1].

Lemma 1. *If \mathbb{A} is a Boolean algebra expansion, then $\tau_{\hat{\mathbb{A}}} = \iota_{\hat{\mathbb{A}}} = \gamma_{\hat{\mathbb{A}}}$.*

Proof. Since $\langle \hat{\mathbb{A}}, \tau_{\hat{\mathbb{A}}} \rangle$ is a compact Hausdorff topological lattice, it follows by [9, Corollary VII-2.3] that $\tau_{\hat{\mathbb{A}}} = \gamma_{\hat{\mathbb{A}}}$. Since $\hat{\mathbb{A}}$ is also a complete, atomic Boolean algebra [4, 19], we know that $\iota_{\hat{\mathbb{A}}} = \gamma_{\hat{\mathbb{A}}}$ [8, Section 2]. \square

3. COMPARING PROFINITE COMPLETION AND MACNEILLE COMPLETION

Theorem 2 (cf. [3, Theorem 4.12]). *Let \mathbb{A} be a modal algebra. TFAE:*

- (1) *the profinite completion $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} , and \diamond is smooth,*
- (2) *\mathbb{A} is residually finite and for every $\theta \in \Phi_{\mathbb{A}}$, $1/\theta$ is a principal lattice filter.*

Proof. If $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} , then $\mu: \mathbb{A} \rightarrow \hat{\mathbb{A}}$ must be injective, so that \mathbb{A} is residually finite (see above). Let $\theta \in \Phi_{\mathbb{A}}$, then it follows from the definition of $\hat{\mathbb{A}}$ that the projection $\pi_{\theta}: \hat{\mathbb{A}} \rightarrow \mathbb{A}/\theta$ commutes with μ and the natural map $a \mapsto a/\theta$; i.e. $a/\theta = \pi_{\theta}\mu(a)$. By [4, Lemma 2.7], $\pi_{\theta}^{-1}(1/\theta)$ is a closed principal filter of $\hat{\mathbb{A}}$; say $\pi_{\theta}^{-1}(1/\theta) = \alpha\uparrow$. Because of the correspondence between modal filters and modal congruences [18, Theorem 29], $\alpha\uparrow$ completely characterizes \mathbb{A}/θ in the following sense: $\mathbb{A}/\theta \cong [0, \alpha]_{\hat{\mathbb{A}}}$ as a bounded lattice [7, Exercise 4.12]. This implies that $\alpha\downarrow$ is finite. Since $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} , $\mu[\mathbb{A}]$ is join-dense in $\hat{\mathbb{A}}$, so by finiteness of $\alpha\downarrow$, there must exist $a \in \mathbb{A}$ such that $\mu(a) = \alpha$. Now $b/\theta = 1/\theta$ iff $\mu(b) \in \pi_{\theta}^{-1}(1/\theta) = \alpha\uparrow = \mu(a)\uparrow$ iff $b \geq a$, so $1/\theta = a\uparrow$ is a principal lattice filter.

Conversely, if \mathbb{A} is residually finite then $\mu: \mathbb{A} \rightarrow \hat{\mathbb{A}}$ is injective, so $(\mu, \hat{\mathbb{A}})$ is a completion of \mathbb{A} . To show that $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} , we will consider the different topologies on $\hat{\mathbb{A}}$. We first show that $\text{At } \hat{\mathbb{A}} \subseteq \mu[\mathbb{A}]$. If $\alpha \in \text{At } \hat{\mathbb{A}}$ there must be some $\theta \in \Phi_{\mathbb{A}}$ such that $\alpha(\theta) \in \text{At } \mathbb{A}/\theta$. Because $1/\theta$ is a principal lattice filter $c\uparrow$, we know that $\mathbb{A}/\theta \cong [0, c]_{\mathbb{A}}$ as a bounded lattice, so there must be some $a \leq c$ with $a \in \text{At } \mathbb{A}$ and $a/\theta = \alpha(\theta)$. But then $\mu(a) = \alpha$. It follows that $\text{At } \hat{\mathbb{A}} \subseteq \mu[\mathbb{A}]$, whence $\iota \subseteq \rho$. Since ι is Hausdorff, so is ρ . Using [17, Theorem 4.5] we can thus conclude that, as far as the Boolean substructure of \mathbb{A} is concerned, $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} . Now to show that \diamond is smooth, remember that $\hat{\diamond}: \hat{\mathbb{A}} \rightarrow \hat{\mathbb{A}}$ is (τ, τ) -continuous. Since $\tau = \iota = \gamma$ by Lemma 1 and $\iota \subseteq \rho$, it follows that $\hat{\diamond}$ is (ρ, γ) -continuous. But then since $\gamma^{\downarrow}, \gamma^{\uparrow} \subseteq \gamma$, it follows by [17, Proposition 5.9] that $\hat{\diamond}^{\bullet} \leq \hat{\diamond} \leq \hat{\diamond}^{\circ}$. Since also $\hat{\diamond}^{\circ} \leq \hat{\diamond}^{\bullet}$ [17, Proposition 5.6], it follows that $\hat{\diamond}$ is smooth. \square

Note that the Theorem above also admits a third equivalent condition, characterizing the dual space of \mathbb{A} . This perspective is further explored in [3].

4. FINITELY GENERATED MODAL ALGEBRAS WITH EDPC

Having equationally definable principal congruences (EDPC) is a strong meta-logical property of varieties of algebras, that coincides with e.g. the existence of a deduction theorem or of a master modality [6, 11] for the modal logic corresponding to a variety of modal algebras. Examples of such logics are logics of bounded depth, n -transitive logics such as **K4**, or regular test-free **PDL** with finitely many basic programs.

Lemma 3 (Proposition 3.4.3 of [11]). *Let \mathcal{V} be a variety of modal algebras. \mathcal{V} has EDPC iff every principal modal filter of an algebra in \mathcal{V} is a principal lattice filter.*

Note that the hypotheses below strongly resemble those of [20, Theorem 4].

Theorem 4 (cf. [3, Corollary 4.5.3]). *If \mathbb{A} is a residually finite, finitely generated modal algebra such that $\text{HSP}(\mathbb{A})$ has equationally definable principal congruences, then the profinite completion $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} and \diamond is smooth.*

Proof. Since \mathbb{A} is a finitely generated algebra, every $\theta \in \Phi_{\mathbb{A}}$ is compact [16, Theorem 1]. As is remarked in [20], every compact congruence of a modal algebra is principal, so under our hypotheses, every $\theta \in \Phi_{\mathbb{A}}$ is principal. Now since $\text{HSP}(\mathbb{A})$ has EDPC,

Lemma 3 tells us that $1/\theta$ is a principal lattice filter for all $\theta \in \Phi_{\mathbb{A}}$, so by Theorem 2, $(\mu, \hat{\mathbb{A}})$ is the MacNeille completion of \mathbb{A} and \diamond is smooth. \square

Note that the EDPC clause in the Theorem above is suppressed in the Heyting algebra case [3], because every variety of Heyting algebras has EDPC.

The conditions of Theorem 4 above are sufficient; what about necessity? From Theorem 2 we know that it is necessary that \mathbb{A} is residually finite. Moreover, in light of [19, Section 3.3], we know that it is necessary for Theorem 4 that \mathbb{A} is atomic. This helps us to find counterexamples to the Theorem if we remove the requirements of being finitely generated or having EDPC. For instance, let \mathbb{A} be the free algebra on 1 generator for the modal logic \mathbf{T} (the logic of reflexive Kripke frames). Then [20, Corollary 7: Example 1] tells us that \mathbb{A} is residually finite and finitely generated but not atomic. This shows us that being residually finite and finitely generated is not sufficient for the conclusion of Theorem 4. Alternatively, the free transitive modal algebra on ω generators is an example of a residually finite modal algebra \mathbb{A} , generating a variety with EDPC, such that \mathbb{A} is not atomic. In summary, residual finiteness is necessary for Theorem 4, and if we remove either of the other two conditions, we can find non-atomic counterexamples.

Corollary 5. *Let \mathbb{A} be a finitely generated free algebra for $\mathbf{K4}$ or \mathbf{PDL} . Then the MacNeille completion and the profinite completion of \mathbb{A} are the same and \diamond is smooth.*

Proof. Let \mathbf{L} be either $\mathbf{K4}$ or \mathbf{PDL} and let \mathcal{V} be the variety corresponding to \mathbf{L} . Since \mathbf{L} has the finite model property, \mathcal{V} is generated by its finite members. By [13, Theorem IV-14.4], this implies that \mathbb{A} , being a finitely generated free algebra for \mathcal{V} , is residually finite. Using the fact that \mathbf{L} has a master modality, it follows that we can apply Theorem 4. \square

REFERENCES

- [1] B. Banaschewski. On profinite universal algebras. In J. Novák, editor, *Proceedings of the Third Prague Topological Symposium*, pages 51–62, September 1971.
- [2] B. Banaschewski and G. Bruns. Categorical Characterization of the MacNeille Completion. *Archiv der Mathematik*, 18:369–377, 1967.
- [3] G. Bezhanishvili and J. Vosmaer. Comparison of MacNeille, canonical and profinite completions. Submitted, 2007.
- [4] Guram Bezhanishvili, Mai Gehrke, Ray Mines, and Patrick J. Morandi. Profinite Completions and Canonical Extensions of Heyting Algebras. *Order*, 23:143–161, 2006.
- [5] Garrett Birkhoff. *Lattice theory*. American Mathematical Society, third edition, 1967.
- [6] Wim Blok and Don Pigozzi. On the structure of varieties with equationally definable principal congruences III. *Algebra Universalis*, 32:545–608, 1994.
- [7] B.A. Davey and H.A. Priestley. *Introduction to Lattices and Order*. Cambridge, second edition, 2002.
- [8] Mai Gehrke and Bjarni Jónsson. Bounded Distributive Lattice Expansions. *Mathematica Scandinavica*, 94:13–45, 2004.
- [9] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, 1980.
- [10] J. Harding. On Profinite Completions and Canonical Extensions. *Algebra Universalis*, 55:293–296, 2006.
- [11] Marcus Kracht. *Tools and Techniques in Modal Logic*. Elsevier, 1999.
- [12] H.M. MacNeille. Partially ordered sets. *Transactions of the American Mathematical Society*, 42:416–160, 1927.
- [13] A.I. Mal'cev. *Algebraic Systems*. Springer-Verlag, 1973.

- [14] H. Rasiowa and R. Sikorski. A Proof of the completeness theorem of Gödel. *Fundamenta Mathematicae*, 37:193–200, 1950.
- [15] Luis Ribes and Pavel Zalesskii. *Profinite Groups*. Springer, 2000.
- [16] Ivan Rival and Bill Sands. A Note on the Congruence Lattice of a Finitely Generated Algebra. *Proceedings of the AMS*, 72:451–455, 1978.
- [17] Mark Theunissen and Yde Venema. MacNeille completions of lattice expansions. To appear in *Algebra Universalis*.
- [18] Yde Venema. Algebras and Coalgebras. In P. Blackburn et al., editors, *Handbook of Modal Logic*. Elsevier, 2007.
- [19] Jacob Vosmaer. Connecting the profinite completion and the canonical extension using duality. Master’s thesis, University of Amsterdam, November 2006.
- [20] F. Wolter. A note on atoms in polymodal algebras. *Algebra Universalis*, 37:334–341, 1997.
E-mail address: `jvosmaer@science.uva.nl`

PLANTAGE MUIDERGRACHT 24, 1018 TV AMSTERDAM, THE NETHERLANDS